



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 57/30, 59/68, 59/84, 51/16, 309/00, 49/76, 211/43	A1	(11) International Publication Number: WO 98/05623 (43) International Publication Date: 12 February 1998 (12.02.98)
(21) International Application Number: PCT/EP97/04050 (22) International Filing Date: 25 July 1997 (25.07.97) (30) Priority Data: MI96A001683 2 August 1996 (02.08.96) IT (71) Applicant (for all designated States except US): DOMPE' S.P.A. [IT/IT]; Via Campo di Pile, I-67100 L'Aquila (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): MANTOVANINI, Marco [IT/IT]; Via Campo di Pile, I-67100 L'Aquila (IT). ALLEGRETTI, Marcello [IT/IT]; Via Campo di Pile, I-67100 L'Aquila (IT). CLAVENNA, Gaetano [IT/IT]; Via Campo di Pile, I-67100 L'Aquila (IT). GANDOLFI, Carmelo [IT/IT]; Via Campo di Pile, I-67100 L'Aquila (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: A PROCESS FOR THE PREPARATION OF 2-ARYL-PROPIONIC AND 2-ARYL-ACETIC ACIDS STARTING FROM ARYL-OLEFINS		
(57) Abstract A process for the preparation of meta-substituted arylalkanoic acids starting from m-aryl-olefins followed by the Claisen rearrangement and an oxidative cleavage of the formed compound.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

A PROCESS FOR THE PREPARATION OF 2-ARYL-PROPIONIC AND 2-ARYL-ACETIC ACIDS STARTING FROM ARYL-OLEFINS

The present invention relates to a process for the preparation of meta-substituted arylalkanoic acids starting from m-aryl-olefins.

A number of meta-substituted arylalkanoic acids are known, being pharmaceutically interesting as analgesic, antipyretic and antiinflammatory agents, such as: 2-amino-3-benzoyl-phenylacetic acid (amfenac), 2-(3-phenoxyphenyl)-propionic acid (fenoprofen), 2-(3-benzoylphenyl)-propionic acid (ketoprofen), 2-(2-hydroxy-5-benzoyl-phenyl)-propionic acid, 2-(2-hydroxy-3-benzoylphenyl)-propionic acid, 2-(2-amino-5-benzoyl-phenyl)-propionic acid, 2-(4-hydroxybenzoyl-phenyl)-propionic acid, 2-(3-hydroxybenzoyl-phenyl)-propionic acid, 3-(2-hydroxybenzoyl-phenyl)-propionic acid, 2-(4-benzoylamino-phenyl)-propionic acid, 2-(3-benzoyl-4-amino-phenyl)-propionic acid, 2-(4-benzoyloxy-phenyl)-propionic acid.

Some of these compounds have a well-established use in therapy; other compounds, such as 2-(2-hydroxy-5-benzoyl-phenyl)-propionic, 2-(2-hydroxy-3-benzoyl-phenyl)-propionic, 2-(4-hydroxy-benzoyl-phenyl)-propionic, 2-(3-hydroxybenzoyl-phenyl)-propionic acids, are examples of non-steroidal antiinflammatories which are metabolites or potential metabolites of ketoprofen.


Ketoprofen is one of the most widely used non-steroidal antiinflammatory agents, with a consumption in the region of hundreds of thousands tons a year. The arrangement of meta-substituents to the aryl-aliphatic residue makes the processes for the total synthesis of

2

these compounds rather expensive and complex, so that up to now no general procedures suitable for their preparation are known.

The available, more established synthetic procedures involve usually reagents (bromine, cyanides, benzene) and/or carboxylation and alkylation reactions which, although formerly usual, are now undesirable in terms of safety of the workers as well as environmental impact, which is nowadays an exceedingly serious problem. X


Now a process has been found for the preparation of meta-substituted arylalkanoic acids of formula



wherein R is hydrogen or C₁-C₆ alkyl, A is a C₁-C₄ alkyl group or aryl, aryloxy or aroyl optionally substituted with one or more alkyl, hydroxy, amino, cyano, nitro, alkoxy, haloalkyl, haloalkoxy substituents;


which comprises:

25 a) Claisen rearrangement of compounds of formula (II):



3

in which A' is C₁-C₄ alkyl group or benzyl, aryl, aryloxy or aroyl optionally substituted with one or more alkyl, hydroxy, amino, cyano, nitro, alkoxy, haloalkoxy substituents, and R is as defined above, X is O, S or NH and the A' group is at the ortho or para position to the X-alkylene group, to give a compound of formula



(III)

10

wherein A' and R are as defined above, XR₁ is at the ortho position to the allyl chain and is a OH, NH₂ or S(C₃-C₁₀) acyl group;

15

b) optional elimination of the -XR₁ group by means of a dehydroxylation, deamination or desulfuration reaction;

c) oxidative cleavage of the compound (III);

20

d) elimination of the -XR₁, OH or NH₂ groups by dehydroxylation or deamination reactions, when not already carried out in step b).

Claisen rearrangement of the compounds of formula II in step a) is effected using conventional conditions, for example using solvents such as hydrocarbons, aromatic hydrocarbons, halo hydrocarbons or mixtures thereof, at a temperature ranging from about 50°C to the solvent's reflux temperature, for times from about 2 to about 10 hours.

30

The compounds of formula II are known or they can be prepared according to known methods, starting from

suitable phenols, anilines or thiophenols, by reaction with suitable allyl or crotyl halides, in the presence of strong bases and of solvents such as halo hydrocarbons, ethers, aromatic hydrocarbons or mixtures thereof.

The OH or NH₂ groups resulting from the Claisen rearrangement at the ortho position to the allyl chain can be removed with conventional methods: in case of amino groups, nitrosation reactions with alkali nitrites will for example be used, followed by decomposition of the diazonium salt in the presence of copper sulfate catalytic amounts. The OH groups, on the other hand, can be removed treating the corresponding trifluoromesylate with formic acid and triethylamine in the presence of palladium acetate/triphenylphosphine complex. Said reactions can be indifferently carried out before or after the double bond degradation reaction to give the carboxylic acid (step c).

The oxidative cleavage of the double bond can be carried out either by means of ozonolysis or using potassium permanganate in phase transfer conditions. During the oxidative cleavage of the double bond in phase transfer conditions with KMnO₄, the benzyl group linked to the aromatic ring is oxidized at the same time into the benzoyl group.

On the other hand, when the XR₁ group is an S-acyl group, known desulfuration reactions can be effected, for example by means of Raney nickel, but such reactions have to be carried out before the oxidative cleavage of the double bond. The desulfuration reaction can be carried out directly on the S-acyl group, or on the SH

5

group, resulting from the hydrolysis of the S-acyl group.

Claisen rearrangement of the compounds of formula II in which X is sulfur takes place in the presence of C₃-C₁₀ fatty acid anhydrides, for example butyric anhydride or higher ones, as described in Tetrahedron Letters 1971, 1969, to give directly the compounds of formula III wherein R₁ is an acyl group having 3 to 10 carbon atoms.

Of course, the synthesis of the compound known as amfenac (see above) requires no elimination of the amino group at the ortho position to the allyl chain.


The process of the invention is particularly advantageous from the environmental point of view, allows to obtain the products in very high yields and provides a versatile synthetic procedure, adaptable to a number of meta-substituted arylacetic or arylpropionic acid derivatives of pharmaceutical interest.

The following examples further illustrate the invention.

EXAMPLE 1

Preparation of Ketoprofen


a) from 4-hydroxy-benzofenone



30 Compound 1 (3 g, 15.1 mmoles) was dissolved in 30 ml of acetone, 2 (4.2 g, 30 mmoles) was added

6

keeping the mixture under stirring. Compound 3 (1.6 ml, 15.5 mmoles) dissolved in 15 ml of acetone was dropped into the mixture, which was kept under stirring at room temperature for 12 hours and subsequently at 40°C for 3 hours. The reaction was complete (TLC control Hexane/EtOAc 8:2). Acetone was evaporated off, the residue was taken up with water (20 ml) and extracted with EtOAc (50 ml). The organic extracts were washed with brine and dried over Na₂SO₄, then filtered and the solvent was evaporated off. 3.8 g (15 mmoles) of the product were recovered (pale yellow oil, solid at + 4°C) sufficiently pure to be used in the subsequent reaction. Hexane/EtOAc 7:3. R_f = 0.38.




2.8 g of compound 4 (11.1 mmoles) were dissolved in DMA (dimethylaniline), heating to a temperature of 210° (temperature of the outer oil bath) for 10 hours (2 x 5 hours). The progress of the reaction was controlled by TLC (Hexane/Etoac 7:3) on a sample taken from the mixture and poured into a 2N HCl solution and extracted with EtOAc.

At the end of 10 hours the starting product had almost disappeared; the mixture was worked up as the control samples. The organic extracts were washed first with 2N HCl to remove completely DMA,

7

then with brine and dried over Na_2SO_4 , filtered and the solvent was evaporated off. A crude weighing 2.55 g was obtained, which was purified on a chromatographic column (Hexane/EtOAc 8:2). The resulting product had still a slightly yellow colour which disappeared upon washing with Hexane:EtOAc 7:3. 1.97 g (7.8 mmoles) of a white solid 5 were obtained. Hexane/EtOAc 7:3. $R_f = 0.14$.

10




15

Compound 5 (1.8 g 7.15 mmoles) was dissolved in 18 ml of dry CH_2Cl_2 under argon atmosphere. The mixture was cooled to a temperature of -25°C with an acetone/dry ice mixture. Compound 7 (1.27 ml, 7.3 mmoles) was added, stirring for 30 minutes, then 6 (1.17 ml, 7.15 mmoles) was added in small portions, checking the temperature. The mixture was kept under stirring at -25°C for a further hour, then it was left to warm at room temperature, diluted with ethyl ether and washed with 2N HCl (3 x 10 ml) and with brine (2 x 10 ml). The mixture was then dried over Na_2SO_4 , filtered and the solvent was evaporated off. 2.5 g of a pale yellow oil 8 were obtained, which was used in the subsequent reaction.

30

Hexane/EtOAc 95:5, $R_f = 0.59$

8




10

Compound 8 (2.5 g, 6.52 mmoles) was dissolved in 18 ml of DMF, 12 (19.4 ml, 2.7 ml) was added at room temperature and subsequently compounds 9 (57 mg, 0.255 mmoles) and 10 (134 mg, 0.511 mmoles) were added under an argon stream. After that, compound 11 (0.6 g, 13.1 moles) was added in small amounts. Temperature was raised to 60°C for 1 hour. The solution changed colour to black. An amount of the mixture was taken with a capillary tube for the control, the content was shaken in 2N HCl and extracted with CH₂Cl₂. The TLC control showed the completion of the reaction. The cooled mixture was poured into 2N HCl and extracted repeatedly with ethyl ether. The organic phase was washed with 2N HCl, sat. NaHCO₃ and brine, then dried over Na₂SO₄ and the solvent was evaporated off. Compound 13 was obtained as a yellow oil (1.45 g, 6.2 mmoles). Hexane/EtOAc 95:5 - R_f = 0.65.

15

20

25



30


9

Compound 13 (1.4 g, 5.93 mmoles) was dissolved in CH_2Cl_2 (35 ml), then water (35 ml), conc. H_2SO_4 (1.89 ml), glacial acetic acid (0.7 ml) and a small amount of Aliquat were added. The mixture was stirred vigorously at room temperature and KMnO_4 (2.7 g, 17 mmoles) was added in small amounts, stirring for 48 hours. A TLC control of the organic phase evidenced the disappearance of the starting product. The mixture was added with $\text{Na}_2\text{S}_2\text{O}_5$ until the brown colour due to manganese dioxide disappears completely, then it was diluted with CH_2Cl_2 and the phases were separated. The organic phase was extracted with sat. NaHCO_3 , the aqueous phase was acidified with 2N HCl and extracted with EtOAc . The organic phase was dried over Na_2SO_4 and evaporated. The crude (950 mg) was taken up into a benzene/petroleum ether 6:20 mixture to crystallize pure Ketoprofen (white solid) (725 mg, 2.85 mmoles). $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5 R_f = 0.14.

EXAMPLE 2


Preparation of Ketoprofen (see example 1)

b) from 4-nitro-benzofenone




10

Compound 15 (13 g, 0.23 moles) was suspended in 50 ml of H₂O, 8 drops of conc. HCl were added and the mixture was heated to ebullition. After 1/2 hour compound 14 (10.4 g, 0.045 moles) dissolved in 50 ml of 95% EtOH was dropped therein. The mixture was heated again to ebullition, keeping reflux for 2 hours. A TLC control (Hexane/EtOAc 7:3) showed the reaction was complete. EtOH was evaporated off, the mixture was alkalized with 2N NaOH, extracted with EtOAc (2x100 ml), dried over Na₂SO₄ and the solvent was evaporated off. A pale yellow solid was obtained, which was washed with an hexane/EtOAc mixture. Compound 16 was obtained as a white solid (7.1 g, 0.036). Hexane/EtOAc 7:3 R_f = 0.3.



Compound 16 (7.1 g, 0.036 moles) was dissolved in dry CH₂Cl₂ (50 ml), 17 (3.46 ml, 0.036 moles) was added at room temperature, stirring for 12 hours. A TLC control (Hexane: EtOAc 7:3) evidenced the disappearance of the starting product. Solvent was evaporated off and the formed acetic acid was removed by evaporation (azeotropic mixture with toluene). 8.3 g (0.035 moles) of 18 as a white solid were obtained. Hexane/EtOAc 7:3 R_f = 0.1.

11




10 Compound 18 (8.2 g, 0.035 moles) was dissolved in hot CH_2Cl_2 (80 ml) (50°C), the solution was left to cool slightly and 19 (2.1 g of a 60% mineral oil dispersion, 0.052 moles) was added. The mixture was heated to 50°C , stirring for 30 minutes. The abundant precipitate formed was added with compound

15 20 (4.23 ml, 0.042 moles) dissolved in 20 ml of dry CH_2Cl_2 . The mixture was kept overnight at room temperature and at 50°C for 8 hours. The starting product had almost disappeared, the mixture was diluted with CH_2Cl_2 , washed with a 5% NaH_2PO_4 solution, then with brine. The organic phase was

20 dried over Na_2SO_4 and the solvent was evaporated off. Compound 21 was obtained as a pale yellow solid, which was sufficiently pure to be used in the subsequent reaction (8.6 g

25 0.029 moles). Hexane/EtOAc 7:3 R_f = 0.44.




30

Compound 21 (8.6 g, 0.029 moles) was dissolved in

12


95% EtOH (15 ml), conc. HCl (15 ml) was added and the mixture was heated to ebullition for 3 hours. The TLC control (Hexane/EtOAc 1:1) evidenced the formation of a UV active spot with $R_f = 0.21$ had not completely disappeared. EtOH was evaporated off, the acid solution was washed with EtOAc to remove the residual starting product, the solution was alkalinized with 5% NaOH and extracted again with EtOAc, dried over Na_2SO_4 , filtered and the solvent was evaporated off. The resulting yellow oil was washed with Hexane/EtOAc 7:3, to precipitate compound 23 as a powdery yellowish solid (5.5 g, 0.022 moles). Hexane/EtOAc 7:3 $R_f = 0.64$.



Compound 23 (0.2 g, 0.8 mmoles) was dissolved in toluene, 24 (0.1 g, 0.8 mmoles) was added and temperature was raised to 100°C for 5 hours. After a night at room temperature, heating was maintained for a further 5 hours. TLC (Hexane/EtOAc 8:2) showed no increases in the desired product. The mixture was left to cool, poured into 2N HCl (5 ml), alkalinized with 2N NaOH and extracted with EtOAc. The extracts were dried over Na_2SO_4 and filtered. The solvent was evaporated off. A crude of about 200 mg was obtained, 150 mg of which were purified on

13

chromatographic column (Hexane: EtOAc 9:1). 18 mg of 25 were obtained (0.07 mmoles). Hexane: EtOAc 8:2 Rf = 0.4



10 Compound 25 (30 mg, 0.12 mmoles) was dissolved in 4 ml of H₂O added with 15 μ l of conc. HCl. The mixture was boiled for 5 minutes, then cooled on ice to 15°C and added with a further 15 μ l of conc. HCl. After stirring for 15 minutes at 10°C, 8.8 mg of 26 (0.127

15 mmoles) dissolved in 0.4 ml of H₂O were added, stirring for 20 minutes. EtOH (3 ml) added with a spatula tip of CuSO₄ was heated to 60°C in another flask. This solution was added with the mixture containing the diazonium salt in small amounts

20 (slight gas evolution), keeping for 30 minutes at 60°C. EtOH was evaporated off, the residue was extracted with EtOAc and dried over Na₂SO₄. The solvent was evaporated off. The crude (30 mg) was purified on preparative TLC Hexane: EtOAc 8:2. 10 mg

25 of compound 28 (pale yellow oil) were obtained. Hexane:EtOAc 8:2 Rf = 0.72.

EXAMPLE 3

4-(perfluoro-1-butanesulfonate)-3-(1'-methyl-2'-propen-1-yl)benzophenone.

30 To a solution of 3-(1'-methyl-2'-propen-1-yl)-4-hydroxybenzophenone (5 g, 0.02 mol) in acetone (25 ml),

14

potassium carbonate (5.5 g, 0.04 mol) and the mixture was left stirring at room temperature for 15'. Then perfluoro-1-butanesulfonyl fluoride (6.6 g, 0.021 mol) was dropped and the solution refluxed for 3 hours. After cooling the inorganic salts were filtered off and the filtrate evaporated under vacuum to give a residue that was diluted with EtOAc (50 ml) and washed with brine (2x50 ml). After drying over Na_2SO_4 the solvent was evaporated to give 4-(perfluoro-1-butanesulfonate)-3-(1'-methyl-2'-propen-1-yl) benzophenone as yellowish oil (10.15 g, 0.019 mol). Yield 95%.

TLC (n-Hexane:EtOAc 9:1) R_f = 0.43.

$^1\text{H-NMR}$ (CDCl_3) δ 7.9-7.82 (dd, 3H, J = 8 Hz), 7.75-7.65 (m, 2H), 7.49 (t, 2H, J = 8Hz), 7.35 (d, 1H, J = 8Hz), 6.05-5.92 (m, 1H), 5.21-5.03 (m, 2H), 3.95 (m, 1H), 1.41 (d, 3H, J = 8Hz).

EXAMPLE 4

3-(1'-methyl-2'-propen-1-yl)benzophenone

To a solution of 4-(perfluoro-1-butanesulfonate)-3-(1'-methyl-2'-propen-1-yl)-benzophenone (6.13 g, 0.0115 mol) in dioxane (50 ml), triethylamine (4.96 ml, 0.036 mol) was added and then, under Ar, palladium acetate (0.108 g, 4.8×10^{-4} mol) and triphenylphosphine (0.25 g, 9.6×10^{-4} mol). Formic acid (1.1 g, 0.024 mol) was then added portionwise and the mixture was warmed at 60°C and left at the temperature for 1.5 h. After cooling the mixture was evaporated under vacuum; the residue was diluted with diethyl ether (50 ml), the formed precipitate was filtered off and the filtrate washed with 2N HCl (2x15 ml), 1N NaOH (2x8 ml), dried over Na_2SO_4 and evaporated under vacuum to give pure 3-(1'-

15

methyl-2'-propen-1-yl)benzophenone (2.6 g 0.011 mol) as yellow oil. Yield =95%

Perfluorobutanesulfonic acid (3.21 g, 0.0107 mol) was recovered from basic aqueous layer (1N NaOH) by acidification (conc. HCl) and extraction with ethyl acetate (2x 20 ml). Recovering yield= 93%.

TLC (n-Hexane/EtOAc 9:1) Rf=0.8

¹H-NMR (CDCl₃) δ 7.79-7.71 (d, 3H), 7.70-7.65 (m, 2H), 7.55-7.35 (m, 4H), 6.09-5.89 (m, 1H), 5.15-5.02 (m, 2H), 3.55 (m, 1H), 1.42 (d, 3H, J= 8Hz).

EXAMPLE 5

4-(benzenesulfonate)-3-(1'-methyl-2'-propen-1-yl)benzophenone.

To a solution of 3-(1'-methyl-2'-propen-1-yl)-4-hydroxybenzophenone (0.9 g, 0.0036 mol) in acetone (10 ml), potassium carbonate (0.99 g, 0.0076 mol) was added and the mixture was left stirring at room temperature for 15'. Then benzenesulfonyl chloride (0.633 g, 0.0036 mol) was dropped and the solution was stirred at room temperature for 3 hours and then refluxed for 1 hour. After cooling inorganic salts were filtered off and the filtrate evaporated under vacuum to give a residue that was diluted with EtOAc (10 ml) and washed with brine (2x10 ml). After drying over Na₂SO₄ the solvent was evaporated to give 4-(benzenesulfonate)-3-(1'-methyl-2'-propen-1-yl)benzophenone as yellowish oil (1.4 g, 0.0036 mol). Quantitative yield.

TLC (n-Hexane:EtOAc 9:1) Rf= 0.4.

¹H-NMR (CDCl₃) δ 7.79-7.71 (dd, 2H, J=8Hz), 7.70-7.4 (m, 10H), 7.25 (d, 1H), 5.9-5.72 (m, 1H), 5.15-4.9 (m, 2H), 3.65 (m, 1H), 1.15 (d, 3H, J= 8Hz).

16

EXAMPLE 63-(1'-methyl-2'-propen-1-yl)benzophenone

To a solution of 4-(benzenesulfonate)-3-(1'-methyl-2'-propen-1-yl)-benzophenone (0.31 g, 0.0008 mol) in DMF (5.5 ml), triethylamine (0.325 mg, 0.0032 mol) was added and then, under Ar, palladium acetate (9×10^{-3} g, 4×10^{-5} mol) and 1,3-Bis(diphenylphosphino)propane (1.8×10^{-4} g, 4.4×10^{-5} mol). Formic acid (0.147 g, 0.0032 mol) was then added portionwise and the mixture was warmed at 90°C and left at the temperature for 3 h. After cooling the mixture was poured in 2N HCl (10 ml) and extracted with EtOAc (2x10 ml). The organic layer was washed with brine (10 ml), dried over Na₂SO₄ and evaporated under vacuum to give a crude oil (0.21 g). Pure 3-(1'-methyl-2'-propen-1-yl)benzophenone (0.085 g 3.5×10^{-4} mol) was obtained as colourless oil by chromatographic purification (silica gel; n-Hexane/EtOAc 8:2). Yield =44%

The saturated compound was isolated in traces as side product.

TLC (n-Hexane/EtOAc 9:1) R_f=0.8

¹H-NMR (CDCl₃) δ 7.79-7.71 (d, 3H), 7.70-7.65 (m, 2H), 7.55-7.35 (m, 4H), 6.09-5.89 (m, 1H), 5.15-5.02 (m, 2H), 3.55 (m, 1H), 1.42 (d, 3H, J = 8Hz).

EXAMPLE 72-(2'-benzenesulfonate-5'-benzoylphenyl)propionic acid

To a solution of 4-(benzenesulfonate)-3-(1'-methyl-2'-propen-1-yl)-benzophenone (1 g, 2.55×10^{-3} mol) in CH₂Cl₂ (15 ml) an equal volume of H₂O, glacial acetic acid (0.3 ml), conc. H₂SO₄ (0.8 ml) and a catalytic

17

amount of Aliquat 336 were added. Then potassium permanganate (1.2 g, 7.6×10^{-3} mol) was added to the mixture portionwise. The mixture was stirred 24 hours at room temperature until disappearance of starting material (TLC). An aqueous solution of sodium metabisulfite was added under stirring until disappearance of the dark brown colour of the solution. CH_2Cl_2 (5 ml) was added and the two phases separated; the organic layer was washed with water (2x 10 ml), dried over Na_2SO_4 and evaporated under vacuum to give a crude residue (1.05 g). The pure 2-(2'-benzenesulfonate-3'-benzoylphenyl)propionic acid (0.8 g, 1.9×10^{-3}) was obtained by chromatographic purification (silica gel, $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5). Yield 76%.

TLC ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 9:1) $R_f=0.45$

$^1\text{H-NMR}$ (CDCl_3) δ 8.0-7.15 (m, 13H), 4.15-3.95 (q, 1H, $J=7\text{Hz}$), 1.35 (d, 3H, $J=7\text{Hz}$).

EXAMPLE 8

2-(2'-perfluorobutanesulfonate-5'-benzoylphenyl)propionic acid

To a solution of 4-(perfluorobutanesulfonate)-3-(1'-methyl-2'-propen-1-yl)benzophenone (0.2 g, 0.37×10^{-3} mol) in CH_2Cl_2 (2.5 ml) an equal volume of H_2O , glacial acetic acid (0.046 ml), conc. H_2SO_4 (0.12 ml) and a catalytic amount of Aliquat 336 were added. Then potassium permanganate (0.173 g, 1.09×10^{-3} mol) was added to the mixture portionwise. The mixture was stirred 24 hours at room temperature until disappearance of starting material (TLC). An aqueous solution of sodium metabisulfite (0.115 g, 1 ml) was added under stirring until disappearance of the dark brown colour of

18

the solution. CH_2Cl_2 (3 ml) was added and the two phases were separated; the organic layer was washed with water (2x 5 ml), dried over Na_2SO_4 and evaporated under vacuum to give a crude residue (0.25 g). 2-(2'-Perfluorobutanesulfonate-5'-benzoylphenyl)propionic acid (0.17 g, 0.3×10^{-3} mol) was obtained by chromatographic purification (silica gel, $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5). Yield 83%. The isolated product was contaminated by traces of Aliquat 336 (4%, NMR).

10 TLC ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 9:1) $R_f=0.5$.

$^1\text{H-NMR}$ (CDCl_3) δ 8.0-7.41 (m, 8H), 4.25-4.1 (q, 1H, $J=7\text{Hz}$), 1.55 (d, 3H, $J=7\text{Hz}$).

EXAMPLE 9

15 3-(1'-methyl-2'-propen-1'-yl)-2-
trifluoromethanesulfonate-diphenylmethane
3-(1'-methyl-2'-propen-1'-yl)-2-(perfluoro-1-
butanesulfonate)-diphenylmethane

20 A solution of crotyl bromide (1, 65 mL) in acetone (18 mL) was added to a stirred mixture of K_2CO_3 (4,5 g) and 2-hydroxydiphenylmethane (2,8 g) in acetone (30 mL). The mixture was kept at room temperature for 8 hrs and then heated for 4 hrs at 45°C .

25 After the usual work-up (filtration of the inorganic material, solvent evaporation and water/ AcOEt extraction) 3,46 g of the 2-crotyloxy-diphenylmethane were recovered.

A solution of 2-crotyloxy-diphenylmethane (2,8 g) in dimethylaniline (30 mL) was warmed for 12 hrs at 210°C (temperature of the outer oil bath).

30 The solvent was removed by vapour stream; the aqueous phase was extracted with AcOEt to give after the

19

usual work-up 2,24 g of 3-(1'-methyl-2'-propen-1'-yl)-2-hydroxy-diphenylmethane that (according to the procedure of the example 1) when treated with trifluoromethanesulfonic anhydride in methylene chloride in the presence of diisopropyl-ethylamine gave 3-(1'-methyl-2'-propen-1'-yl)-2-trifluoromethanesulfonate-diphenylmethane. By reaction with perfluoro-1-butanefluoride (according to the procedure of the example 3), the same product gave 3-(1'-methyl-2'-propen-1'-yl)-2-(perfluoro-1-butanefluoride)-diphenylmethane.

EXAMPLE 10

4-(perfluoro-1-butanefluoride)-3-(1'-methyl-2'-propen-1'-yl)-diphenylmethane

The use of the 4-hydroxydiphenylmethane [obtained by Paternò reaction between phenol and benzyl alcohol; $^1\text{H-NMR}$ δ 7.3-6.5 (m, 9H), 4.9 (s, OH), 3.9 (s, 2H) TLC (n-Hexane/EtOAc 8:2 R_f =0.50)] in the procedure of the example 9 instead of commercially available 2-hydroxydiphenylmethane [$^1\text{H-NMR}$ δ 7.3-6.5 (m, 9H), 4.65 (s, OH), 3.95 (s, 2H) TLC (n-Hexane/EtOAc 8:2 R_f =0.55)] gave the 4-crotyloxy-diphenylmethane.

By Claisen rearrangement of 3,2 g of 4-crotyloxy-diphenylmethane in the dimethylaniline 2,87 g of 3-(1'-methyl-2'-propen-1'-yl)-4-hydroxy-diphenylmethane were obtained. In the presence of finely powdered K_2CO_3 (3g), a solution of perfluoro-1-butanefluoride (3,3 g) in acetone (6mL) was dropwise added to a solution of 2,5 g of this material in acetone (10 ml).

The mixture was refluxed for 3 hrs, the inorganic salts were filtered off to obtain after the usual work-

20

up 5,3 g of 4-(perfluoro-1-butanesulfonate)-3-(1'-methyl-2'-propen-1'-yl)-diphenylmethane.

EXAMPLE 11

Using in the procedure of example 8 a
5 diphenylmethane-sulfonate consisting of:

3-(1'-methyl-2'-propen-1'-yl)-2-trifluoromethanesulfonate-diphenylmethane, 3-(1'-methyl-2'-propen-1'-yl)-4-(perfluoro-1-butanesulfonate)-diphenylmethane;

3-(1'-methyl-2'-propen-1'-yl)-2-(perfluoro-1-butanesulfonate)-diphenylmethane;

after potassium permanganate phase transfer oxidation, the following acids:

2-(2'-trifluoromethanesulfonate-3'-benzoylphenyl)-propionic acid;

15 2-(2'-perfluorobutanesulfonate-3'-benzoylphenyl)-propionic acid;

2-(2'-perfluorobutanesulfonate-5'-benzoylphenyl)-propionic acid were obtained.

Using the above sulfonates in the procedure of the
20 examples 1, 4, and 6, by reductive removal of the sulfonate group with formic acid in the presence of Pd acetate and a suitable phosphine (triphenylphosphine and 1,3-bis-(diphenyl-phosphine)propane], the same 2-(3-benzoylphenyl)-propionic acid (ketoprofen) was obtained.

25 EXAMPLE 12

Methyl esters of 2-(2'-benzenesulfonate-3'-benzoylphenyl)propionic acid and 2-(2'-perfluorobutanesulfonate-3'-benzoylphenyl) propionic acid were easily
30 obtained starting from the acids according to the usual esterification methods (diazomethane/ether; dry MeOH/ and catalytic amounts of H₂SO₄, benzenesulfonic or p-

toluensulfonic acids).

By hydrogenolysis of the following esters:

2-(2'-perfluorobutane sulfonate-5'-benzoylphenyl) propionic acid methyl ester.

5 $^1\text{H-NMR}$ (CDCl_3) δ 7.8-7.4 (m, 8H), 4.25-4.1 (q, 1H, $J=7\text{Hz}$), 3.95 (s, 3H) 1.55 (d, 3H, $J=7\text{Hz}$).

2-(2'-benzenesulfonate-5'-benzoylphenyl)propionic acid methyl ester.

10 $^1\text{H-NMR}$ (CDCl_3) δ 8.1-7.9 (d, 1H), 7.9-7.5 (m, 11H), 7.33 (d, 1H $J=8\text{Hz}$), 4.15-3.95 (q, 1H, $J=7\text{Hz}$), 3.75 (s, 3H), 1.35 (d, 3H, $J=7\text{Hz}$).

The 2-(3'-benzoylphenyl)propionic acid methyl ester (methyl ketoprofenate) was obtained according the following procedure.

15 To a solution of 2-(2'-perfluorobutane sulfonate - 5'-benzoylphenyl) propionic acid methyl ester (6.8 g, 0.012 mol) in dioxane (50 ml), triethylamine (4.96 mL, 0.036 mol) was added and then, under Ar, palladium acetate (0.108 g, 4.8×10^{-4} mol) and
20 triphenylphosphine (0.25 g, 9.6×10^{-4} mol). Formic acid (1.1 g, 0.024 mol) was then added portionwise and the mixture was warmed at $T=60^\circ\text{C}$ for 1.5 h. After cooling the mixture was evaporated under vacuum; the residue was diluted with diethyl ether (50 ml), the
25 formed precipitate was filtered off. The filtrate was washed with 2N HCl (2x15 ml), 1N NaOH (2x8 mL), dried over Na_2SO_4 and evaporated under vacuum to give pure 2-(3'-benzoylphenyl)propionic acid methyl ester (2.14 g 0.008 mol) as yellow oil. Yield =67%; TLC (n-Hexane/EtOAc 9:1) $R_f=0.53$. $^1\text{H-NMR}$ (CDCl_3) δ 7.85-7.3 (m, 9H), 3.85 (q, 1H, $J=8\text{Hz}$), 3.65 (s, 3H), 1.5 (d, 3H, $J=$
30

8Hz).

Perfluoro-1-butansulfonic acid (3.21 g, 0.0107 mol) was recovered from basic aqueous layer (NaOH 1N) by acidification (HCl conc.) and extraction with ethyl acetate (2x 20 ml).

EXAMPLE 13

3-(1'-methyl-2'-propen-1-yl)-2-aminodiphenylmethane

3-(1'-methyl-2'-propen-1-yl)-4-aminodiphenylmethane

Using in the procedure of the example 2 both 2-amino-diphenylmethane and 4-amino-diphenylmethane, the following amines

3-(1'-methyl-2'-propen-1-yl)-2-aminodiphenylmethane
¹H-NMR δ 7.5-7.2 (m, 5H), 7.17 (d, 1H J= 9 Hz), 7.0 (d, 1H J= 9Hz); 6.85 (t, 1H J=9Hz); 6.1-5.9 (m, 1H); 5.2-5.0 (m, 2H); 3.96 (s, 2H); 3.6 (bs, NH₂); 3.5 (m, 1H), 1.35 (d, 3H, J= 7 Hz) TLC (n-Hexane/EtOAc 9:1 R_f=0.2).

3-(1'-methyl-2'-propen-1-yl)-4-aminodiphenylmethane
¹H-NMR δ 7.5-7.1 (m, 5H), 6.85-6.7 (m, 3H); 6.1-5.9 (m, 1H); 5.2-5.0 (m, 2H); 4.3 (s, 2H); 4.2 (bs, NH₂); 3.5 (m, 1H), 1.35 (d, 3H, J= 7 Hz) TLC (n-Hexane/EtOAc 9:1) were obtained.

EXAMPLE 14

3-(1'-methyl-2'-propen-1-yl)-2-hydroxy-diphenylmethane

3-(1'-methyl-2'-propen-1-yl)-4-hydroxy-diphenylmethane

A mixture of 3-(1'-methyl-2'-propen-1-yl)-2-aminodiphenylmethane (0.3 g, 1.26 mmol), water (3 ml) and conc HCl (0.114 ml 1.38 mmol.) was heated at the reflux temperature for 5 min. After cooling at room temperature, an equal amount of conc. HCl was added to the mixture, that is cooled at 4 °C and treated with an aqueous solution of sodium nitrite (0.096 g,

23

1.38 mmol. in 1 mL), maintaining the temperature of the mixture below 5 °C. The subsequent addition of hypophosphorous acid (1.3 ml 50% sol., 12.6 mmol) gave 3-(1'-methyl-2'-propen-1-yl)-2-hydroxy-diphenylmethane.

¹H-NMR δ 7.4-7.15 (m, 5H), 7.1-6.95 (m, 2H), 6.9-6.8 (m, 1H); 6.2-6.0 (m, 1H); 5.2-5.0 (m, 2H); 4.1 (s, 2H); 3.8 (m, 1H); 1.35 (d, 3H, J= 7 Hz) TLC (n-Hexane/EtOAc 9:1 R_f=0.39).

Starting from the 4-amine-diphenyl-methane-propenyl derivate, the 3-(1'-methyl- 2'-propen-1-yl)-4-hydroxy-diphenylmethane was prepared.

EXAMPLE 15

Starting from 3-(1'-methyl-2'-propen-1-yl)-2-amino-diphenylmethane and making use of well-known acylation procedures 3-(1'-methyl-2'-propen-1-yl)-2-acetylamino-diphenylmethane and 3-(1'-methyl-2'-propen-1-yl)-2-tert-butoxycarbonylamino-diphenyl methane are prepared.


According the procedure of example 1, the following potassium permanganate phase transfer oxidation of the above N-acylamino compounds affords respectively: 2-(2'-acetylamino-3'-benzoylphenyl)-propionic acid, 2-(2'-tert-butoxycarbonylamino-3'-benzoylphenyl)-propionic acid.

The treatment of 2-(2'-tert-butoxycarbonylamino-3'-benzoylphenyl)-propionic acid in CH₂Cl₂, at 5-10 °C, with an excess of trifluoroacetic acid allows to recover the 2-(2-amino-3-benzoylphenyl)propionic acid as trifluoroacetic acid salt.

30

CLAIMS

1. A process for the preparation of meta-substituted arylalkanoic acids of formula




10 wherein R is hydrogen or C₁-C₆ alkyl, A is a C₁-C₄ alkyl group or aryl, aryloxy or aroyl optionally substituted with one or more alkyl, hydroxy, amino, cyano, nitro, alkoxy, haloalkyl,

15 haloalkoxy substituents;

which comprises:


a) Claisen rearrangement of compounds of formula (II):



25 in which A' is C₁-C₄ alkyl group or benzyl, aryl, aryloxy or aroyl optionally substituted with one or more alkyl, hydroxy, amino, cyano, nitro, alkoxy, haloalkoxy substituents and R is as defined above, X is O, S or NH and the A' group is at the ortho or para position to the X-

30 alkylene group, to give a compound of formula

25



(III)

5

wherein A' and R are as defined above, XR₁ is at the ortho position to the allyl chain and is a OH, NH₂ or S(C₃-C₁₀) acyl group;

10

b) optional elimination of the -XR₁ group by means of a dehydroxylation, deamination or desulfuration reaction;

c) oxidative cleavage of the compound (III);

d) elimination of the -XR₁, OH or NH₂ groups by dehydroxylation or deamination reactions, when not

15

already carried out in step b)).

2. A process according to claim 1, for the preparation of the compound of formula I in which R is methyl and A is benzoyl.

20

3. A process according to claim 1 or 2, wherein compounds of formula II in which X is O are used.

4. A process according to claim 1 or 2, wherein compounds of formula II in which X is NH are used.

5. A process according to claim 1 or 2, wherein compounds of formula II in which X is S are used.

25

6. A process according to any one of the above claims, wherein step c) is effected with KMnO₄ in phase transfer conditions.

7. Compounds of formula II wherein R, X and A are as defined above.

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/EP 97/04050

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C57/30 C07C59/68 C07C59/84 C07C51/16 C07C309/00
C07C49/76 C07C211/43

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 025 397 A (CHINOIN GYOGYSZER ES VEGYESZETI TERMEKEK GYARA RT) 23 January 1980 ENTIRE DOCUMENT	1-3,6,7
Y	GB 1 586 798 A (CHINOIN GYOGYSZER ES VEGYESZETI TERMEKEK GYARA RT) 25 March 1981 see claims 1,2	1
Y	EP 0 394 949 A (MITSUBISHI PETROCHEMICAL CO.) 31 October 1990 see claims 1-4	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 December 1997

Date of mailing of the international search report

16/12/1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Klag, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/04050

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2025397 A	23-01-80	AT 367013 B	25-05-82
		CA 1122603 A	27-04-82
		CH 646133 A	15-11-84
		DK 209879 A	24-11-79
		FR 2438026 A	30-04-80
		JP 55017369 A	06-02-80
		NL 7904056 A	27-11-79
		SE 7904396 A	24-11-79
		US 4317920 A	02-03-82
		US 4385004 A	24-05-83
		US 4465855 A	14-08-84
GB 1586798 A	25-03-81	CH 627727 A	29-01-82
		DK 517977 A	24-05-78
		JP 53082751 A	21-07-78
		NL 7712841 A	25-05-78
		SE 7713129 A	24-05-78
		SU 895282 A	30-12-81
EP 0394949 A	31-10-90	JP 2282343 A	19-11-90
		JP 3066638 A	22-03-91
		US 5087766 A	11-02-92